


Case report

V. de Lédinghen
CHU Bordeaux



John, 45 year-old

- 1998 : diagnosis of HCV infection
- Genotype 1b
- Contamination : transfusion 1985
- Excessive alcohol use until 1994
- 1.83 m ; 85 kg
- Liver biopsy : A2F2
- Treatment: interferon and ribavirin
- No response (positive RNA at W12)
- Stop treatment
- Lost of follow-up

-
- 2003 : new visit
 - No alcohol or tobacco use
 - Viral load 950 000 IU/ml
 - Platelets 150 G/L
 - PT 93 %
 - ALT 115 IU/l
 - AST 95 IU/l

Non-invasive evaluation of liver fibrosis

FibroScan 27.6 +/- 8.1 kPa  F4

Fibrotest ALT 89 IU/ml
 GGT 81 IU/l
 A2M 2,8 g/l
 Bili 20 µmol/l
 Haptoglobin 0.41 g/l
 APOA1 1.48 g/l  F2

0,56

What do you think ?

1. FibroScan is correct
2. Fibrotest is correct
3. I don't know

What do you do ?

1. I repeat FibroScan and Fibrotest 6 months later
2. I use other non-invasive tests
3. I perform a liver biopsy

6 months later

FibroScan $17.5 +/ - 2.3 \text{ kPa}$  F4

Fibrotest ALT 58 IU/l
 GGT 60 IU/l
 Bili 14
 A2M 2.7  F2
 Haptoglobin 0.71
 APOA1.11
 0,58

What do you do ?

1. I don't need liver fibrosis evaluation: I treat!
2. I perform a liver biopsy
3. I wait 6 months and repeat evaluation

Liver biopsy: 15 mm A2F4

- Endoscopy: no oesophageal varices
- US: non hepatocellular carcinoma
- Treatment: Pegasys 180 µg/w + Ribavirin 1,200 mg/d
- W4 100 000 IU/ml
- Hemoglobin 11 g/dl
- Platelets 80 G/l

Treatment 48 weeks

- Negative HCV RNA at W12 and W48
- Normal ALT level

Do you perform NI evaluation of liver fibrosis at week 48?

1. Yes
2. No

At week 48

- Platelets: 99 G/L
- ALT 42 IU/L
- FibroScan : 14.6 kPa (IQR 4.8)
- Fibrotest: 0.53

Do you perform NI evaluation of liver fibrosis at week 72?

1. Yes
2. No

Week 72

- Negative HCV RNA
- ALT 38 IU/L
- GGT 62 IU/L
- Platelets: 176 G/L
- FibroScan 9.9 kPa (IQR 0.7)
- Fibrotest 0.56

What do you think ?

1. There is always a cirrhosis
2. Fibrosis could be F3
3. Fibrosis could be F2

Follow-up

	2005	2006	2007
ALT	38	30	44
HCV RNA	-	-	-
FibroScan (kPa)	9.9	8.6	7.2
FibroTest	0.56	0.55	0.57

What do you think ?

1. There is always a cirrhosis
2. Fibrosis is F2 or F3
3. I don't know

Liver biopsy

- A1 F3/F4

Conclusion

- Without liver biopsy
- Even if there is a regression of liver stiffness

Cirrhosis once, cirrhosis forever!